Characteristics of Stars

Reading Preview Key Concepts

- How are stars classified?
- How do astronomers measure distances to the stars?
- What is an H-R diagram and how do astronomers use it?

Key Terms

- constellation
- spectrograph
- apparent brightness
- absolute brightness
- light-year
- parallax
- Hertzsprung-Russell diagram
- main sequence
- Using Prior Knowledge Before you read, write what you know about the characteristics of stars in a graphic organizer like the one below. As you read, write what you learn.

What You Know 1. Stars are bright and hot. 2.

What You Learned 1. 2.


Lah Discover Activity

How Does Your Thumb Move?

- 1. Stand facing a wall, at least an arm's length away. Stretch your arm out with your thumb up and your fingers curled.
- 2. Close your right eye and look at your thumb with your left eye. Line your thumb up with something on the wall.
- 3. Now close your left eye and open your right eye. How does your thumb appear to move along the wall?
- 4. Bring your thumb closer to your eye, about half the distance as before. Repeat Steps 2 and 3.

Think It Over

Observing How does your thumb appear to move in Step 4 compared to Step 3? How are these observations related to how far away your thumb is at each step? How could you use this method to estimate distances?



Different cultures gave different names to the constellations. For example, a large constellation in the winter sky is named Orion, the Hunter, after a Greek myth. In this constellation, Orion is seen with a sword in his belt and an upraised arm. The ancient Sumerians thought that the stars in Orion formed the outline of a sheep. In ancient China, this group of stars was called "three," probably because of the three bright stars in Orion's belt.

Astronomers use the patterns of the constellations to locate objects in the night sky. But although the stars in a constellation look as if they are close to one another, they generally are not. They just happen to lie in the same part of the sky as seen from Earth.

Illustration of Orion ▼



Classifying Stars

Like the sun, all stars are huge spheres of glowing gas. They are made up mostly of hydrogen, and they produce energy through the process of nuclear fusion. This energy makes stars shine brightly. Astronomers classify stars according to their physical characteristics. Characteristics used to classify stars include color, temperature, size, composition, and brightness.

Color and Temperature If you look at the night sky, you can see slight differences in the colors of the stars. For example, Betelgeuse (BAY tul jooz), the bright star in Orion's shoulder, looks reddish. Rigel, the star in Orion's heel, is blue-white.

Like hot objects on Earth, a star's color reveals its surface temperature. If you watch a toaster heat up, you can see the wires glow red-hot. The wires inside a light bulb are even hotter and glow white. Similarly, the coolest stars—with a surface temperature of about 3,200 degrees Celsius—appear reddish in the sky. With a surface temperature of about 5,500 degrees Celsius, the sun appears yellow. The hottest stars in the sky, with surface temperatures of over 20,000 degrees Celsius, appear bluish.

Size When you look at stars in the sky, they all appear to be points of light of the same size. Many stars are actually about the size of the sun, which is a medium-sized star. However, some stars are much larger than the sun. Very large stars are called giant stars or supergiant stars. If the supergiant star Betelgeuse were located where our sun is, it would be large enough to fill the solar system as far out as Jupiter.

Most stars are much smaller than the sun. White dwarf stars are about the size of Earth. Neutron stars are even smaller, only about 20 kilometers in diameter.



For: More on types of stars Visit: PHSchool.com Web Code: cfd-5042

FIGURE 5 Star Size

Stars vary greatly in size. Giant stars are typically 10 to 100 times larger than the sun and more than 1,000 times the size of a white dwarf. Calculating Betelgeuse has a diameter of 420 million kilometers. How many times larger is this than the sun, which has a diameter of 1.4 million kilometers?





FIGURE 6
Spectrums of Four Stars

Astronomers can use line spectrums to identify the chemical elements in a star. Each element produces a characteristic pattern of spectral lines.



Chemical Composition Stars vary in their chemical composition. The chemical composition of most stars is about 73 percent hydrogen, 25 percent helium, and 2 percent other elements by mass. This is similar to the composition of the sun.

Astronomers use spectrographs to determine the elements found in stars. A **spectrograph** (SPEK truh graf) is a device that breaks light into colors and produces an image of the resulting spectrum. Most large telescopes have spectrographs.

The gases in a star's atmosphere absorb some wavelengths of light produced within the star. When the star's light is seen through a spectrograph, each absorbed wavelength is shown as a dark line on a spectrum. Each chemical element absorbs light at particular wavelengths. Just as each person has a unique set of fingerprints, each element has a unique set of lines for a given temperature. Figure 6 shows the spectral lines of four elements. By comparing a star's spectrum with the spectrums of known elements, astronomers can infer how much of each element is found in the star.




What is a spectrograph?

Lab zone Skills Activity

Inferring

The lines on the spectrums below are from three different stars. Each of these star spectrums is made up of an overlap of spectrums from the individual elements shown in Figure 6. In star A, which elements have the strongest lines? Which are the strongest in star B? In star C?



Brightness of Stars

Stars also differ in brightness, the amount of light they give off. The brightness of a star depends upon both its size and temperature. Recall that the photosphere is the layer of a star that gives off light. Betelgeuse is fairly cool, so a square meter of its photosphere doesn't give off much light. But Betelgeuse is very large, so it shines brightly.

Rigel, on the other hand, is very hot, so each square meter of Rigel's photosphere gives off a lot of light. Even though it is smaller than Betelgeuse, Rigel shines more brightly. How bright a star looks from Earth depends on both its distance from Earth and how bright the star truly is. Because of these two factors, the brightness of a star can be described in two ways: apparent brightness and absolute brightness.

Apparent Brightness A star's apparent brightness is its brightness as seen from Earth. Astronomers can measure apparent brightness fairly easily using electronic devices. However, astronomers can't tell how much light a star gives off just from the star's apparent brightness. Just as a flashlight looks brighter the closer it is to you, a star looks brighter the closer it is to Earth. For example, the sun looks very bright. This does not mean that the sun gives off more light than all other stars. The sun looks so bright simply because it is so close. In reality, the sun is a star of only average brightness.

Absolute Brightness A star's absolute brightness is the brightness the star would have if it were at a standard distance from Earth. Finding a star's absolute brightness is more complex than finding its apparent brightness. An astronomer must first find out both the star's apparent brightness and its distance from Earth. The astronomer can then calculate the star's absolute brightness.

Astronomers have found that the absolute brightness of stars can vary tremendously. The brightest stars are more than a billion times brighter than the dimmest stars!



What is a star's absolute brightness?



Star Bright

You can compare absolute and apparent brightness.

- 1. Dim the lights. Put two equally bright flashlights next to each other on a table. Turn them on.
- 2. Look at the flashlights from the other side of the room. Think of the flashlights as two stars. Then compare them in terms of absolute and apparent brightness.
- 3. Move one of the flashlights closer to you and repeat Step 2.
- 4. Replace one of the flashlights with a brighter one. Repeat Steps 1 and 2 with the unequally bright flashlights.

Making Models How could you place the flashlights in Step 4 so that they have the same apparent brightness? Try it.



Absolute Brightness
The streetlights in this photo all give off about the same amount of light, and so have about the same absolute brightness.

Applying Concepts Why do the closer streetlights appear brighter than the more distant lights?

Measuring Distances to Stars

Imagine that you could travel to the stars at the speed of light. To travel from Earth to the sun would take about 8 minutes, not very much time for such a long trip. The next nearest star, Proxima Centauri, is much farther away. A trip to Proxima Centauri at the speed of light would take 4.2 years!

The Light-Year Distances on Earth's surface are often measured in kilometers. However, distances to the stars are so large that kilometers are not very practical units. Astronomers use a unit called the light-year to measure distances between the stars. In space, light travels at a speed of about 300,000 kilometers per second. A light-year is the distance that light travels in one year, about 9.5 million million kilometers.

Note that the light-year is a unit of distance, not time. To help you understand this, consider an everyday example. If you bicycle at 10 kilometers per hour, it would take you 1 hour to go to a mall 10 kilometers away. You could say that the mall is "1 bicycle-hour" away.

Parallax Standing on Earth looking up at the sky, it may seem as if there is no way to tell how far away the stars are. However, astronomers have found ways to measure those distances. Astronomers often use parallax to measure distances to nearby stars.


Parallax is the apparent change in position of an object when you look at it from different places. For example, imagine that you and a friend have gone to a movie. A woman with a large hat sits down in front of you, as shown in Figure 8. Because you and your friend are sitting in different places, the woman's hat blocks different parts of the screen. If you are sitting on her left, the woman's hat appears to be in front of the large dinosaur. But to your friend on the right, she appears to be in front of the bird.

Have the woman and her hat moved? No. But because you changed your position, she appears to have moved. This apparent movement when you look from two different directions is parallax.

Parallax at the Movies
You and your friend are sitting behind a woman with a large hat. Applying Concepts Why is your view of the screen different from your friend's view?








Parallax in Astronomy Astronomers are able to measure the parallax of nearby stars to determine their distances. As shown in Figure 9, astronomers look at a nearby star when Earth is on one side of the sun. Then they look at the same star again six months later, when Earth is on the opposite side of the sun. Astronomers measure how much the nearby star appears to move against a background of stars that are much farther away. They can then use this measurement to calculate the distance to the nearby star. The less the nearby star appears to move, the farther away it is.

Astronomers can use parallax to measure distances up to a few hundred light-years from Earth. The parallax of any star that is farther away is too small to measure accurately.



How is parallax useful in astronomy?

FIGURE 9 Parallax of Stars

The apparent movement of a star when seen from a different position is called parallax. Astronomers use parallax to calculate the distance to nearby stars. Note that the diagram is not to scale.

Interpreting Diagrams Why do nearby stars appear to change position between January and July?

The Hertzsprung-Russell Diagram

About 100 years ago, two scientists working independently made the same discovery. Both Ejnar Hertzsprung (EYE nahr HURT sprung) in Denmark and Henry Norris Russell in the United States made graphs to find out if the temperature and the absolute brightness of stars are related. They plotted the surface temperatures of stars on the x-axis and their absolute brightness on the y-axis. The points formed a pattern. The graph they made is still used by astronomers today. It is called the Hertzsprung-Russell diagram, or H-R diagram.





FIGURE 10
The Hertzsprung-Russell diagram shows the relationship between the surface temperature and absolute brightness of stars.
Interpreting Diagrams Which star has a hotter surface: Rigel or Aldebaran?

Astronomers use H-R diagrams to classify stars and to understand how stars change over time. As you can see in Figure 10, most of the stars in the H-R diagram form a diagonal area called the main sequence. More than 90 percent of all stars, including the sun, are main-sequence stars. Within the main sequence, surface temperature increases as absolute brightness increases. Thus, hot bluish stars are located at the left of an H-R diagram and cooler reddish stars are located at the right of the diagram.

The brightest stars are located near the top of an H-R diagram, while the dimmest stars are located at the bottom. Giant and supergiant stars are very bright. They can be found near the top center and right of the diagram. White dwarfs are hot, but not very bright, so they appear at the bottom left or bottom center of the diagram.



What is the main sequence?




Figure 11
Orion
Orion includes the red supergiant
Betelgeuse, the blue supergiant Rigel, and
many other main-sequence and giant stars.

Section 2 Assessment

Target Reading Skill Using Prior Knowledge Review your graphic organizer and revise it based on what you just learned in the section.

Reviewing Key Concepts

- 1. a. Listing Name three characteristics used to classify stars.
 - b. Comparing and Contrasting What is the difference between apparent brightness and absolute brightness?
 - c. Applying Concepts Stars A and B have about the same apparent brightness, but Star A is about twice as far from Earth as Star B. Which star has the greater absolute brightness? Explain your answer.
- 2. a. Measuring What is a light-year?
 - b. Defining What is parallax?
 - c. Predicting Vega is 25.3 light-years from Earth and Arcturus is 36.7 light-years away. Which star would have a greater parallax? Explain.

- 3. a. Summarizing What two characteristics of stars are shown in an H-R diagram?
 - b. Identifying Identify two ways in which astronomers can use an H-R diagram.
 - c. Classifying The star Procyon B has a surface temperature of 6,600°C and an absolute brightness that is much less than the sun's. What type of star is Procyon B? (Hint: Refer to the H-R diagram.)

Zone At-Home Activity

Observing Orion With adult family members, go outside on a clear, dark night. Determine which way is south. Using the star charts in the appendix, look for the constellation Orion, which is visible in the evening during winter and spring. Find the stars Betelgeuse and Rigel in Orion and explain to your family why they are different colors.

Skills Lab

How Far Is That Star?

Problem

How can parallax be used to determine distances?

Skills Focus

inferring, calculating, predicting

Materials

- masking tape
 paper clips
 pen
- black and red pencils
 metric ruler
 paper
- meter stickcalculator
- lamp without a shade, with 100-watt light bulb
- copier paper box (without the lid)
- flat rectangular table, about 1 m wide

Procedure 🕅




PARM Telescope Model

- 1. Place the lamp on a table in the middle of the classroom.
- 2. Carefully use the tip of the pen to make a small hole in the middle of one end of the box. The box represents a telescope.

- 3. At the front of the classroom, place the box on a flat table so the hole points toward the lamp. Line the left side of the box up with the left edge of the table.
- 4. Put a small piece of tape on the table below the hole. Use the pen to make a mark on the tape directly below the hole. The mark represents the position of the telescope when Earth is on one side of its orbit

PART Star 1

- 5. Label a sheet of paper Star 1 and place it inside the box as shown in the drawing. Hold the paper in place with two paper clips. The paper represents the film in a telescope.
- 6. Darken the room. Turn on the light to represent the star.
- 7. With the red pencil, mark the paper where you see a dot of light. Label this dot A. Dot A represents the image of the star on the film.
- 8. Move the box so the right edge of the box lines up with the right edge of the table. Repeat Step 4. The mark on the tape represents the position of the telescope six months later, when Earth is on the other side of its orbit.



Data Table							
Star	Parallax Shift (mm)	Focal Length (mm)	Diameter of Orbit (mm)	Calculated Distance to Star (mm)	Calculated Distance to Star (m)	Actual Distance to Star (m)	

- Repeat Step 7, using a black pencil to mark the second dot B. Dot B represents the image of the star as seen 6 months later from the other side of Earth's orbit.
- 10. Remove the paper. Before you continue, copy the data table into your notebook.
- 11. Measure and record the distance in millimeters between dots A and B. This distance represents the parallax shift for Star 1.
- 12. Measure and record the distance from the hole in the box to the lamp. This distance represents the actual distance to the star.
- 13. Measure and record the distance from the hole (lens) to the back of the box in millimeters. This distance represents the focal length of your telescope.
- 14. Measure and record the distance in millimeters between the marks on the two pieces of masking tape. This distance represents the diameter of Earth's orbit.

PARE Stars 2 and 3

- 15. Move the lamp away from the table—about half the distance to the back of the room. The bulb now represents Star 2. Predict what you think will happen to the light images on your paper.
- 16. Repeat Steps 6-12 with a new sheet of paper to find the parallax shift for Star 2.
- 17. Move the lamp to the back of the classroom. The bulb now represents Star 3. Repeat Steps 6–12 with a new sheet of paper to find the parallax shift for Star 3.

Analyze and Conclude

- 1. Inferring What caused the apparent change in position of the dots of light for each star? Explain.
- 2. Calculating Use the following formula to calculate the distance from the telescope to Star 1.

Distance =
$$\frac{\text{Diameter} \times \text{Focal length}}{\text{Parallax shift}}$$

- 3. Calculating Divide your result from Question 2 by 1,000 to get the distance to the light bulb in meters.
- 4. Calculating Repeat Questions 2 and 3 for Stars 2 and 3.
- 5. Predicting Was your prediction in Step 15 correct? Why or why not?
- 6. Interpreting Data How did your calculation for Star 3 compare with the actual distance? What could you do to improve your results?
- 7. Communicating Write a paragraph that explains how parallax shift varies with distance. Relate each star's parallax shift to its distance from Earth.

Design an Experiment

What would happen if you kept moving the lamp away from the box? Is there a distance at which you can no longer find the distance to the star? Design an experiment to find out.

Lives of Stars

Reading Preview

Key Concepts

- How does a star form?
- What determines how long a star will exist?
- What happens to a star when it runs out of fuel?

Key Terms

- nebula protostar
- white dwarf supernova
- neutron star pulsar
- black hole

ঠি) Target Reading Skill

Sequencing As you read, make a flowchart like the one below that shows the stages in the life of a star like the sun. Write each step of the process in a separate box in the flowchart in the order that it occurs.

Life Cycle of a Sun-like Star


Protostar forms from a nebula.

A star is born as fusion begins.

Lab Discover Activity

What Determines How Long Stars Live?

- This graph shows how the mass of a star is related to its lifetime how long the star lives before it runs out of fuel.
- 2. How long does a star with 0.75 times the mass of the sun live? How long does a star with 3 times the mass of the sun live?



Think It Over

Drawing Conclusions Describe the general relationship between a star's mass and its lifetime.

Imagine that you want to study how people age. You wish you could watch a few people for 50 years, but your project is due next week! You have to study a lot of people for a short time, and classify the people into different age groups. You may come up with groups like *babies*, young adults, and elderly people. You don't have time to see a single person go through all these stages, but you know the stages exist.

Astronomers have a similar problem in trying to understand how stars age. They can't watch a single star for billions of years. Instead, they study many stars and other objects in

space. Over time, astronomers have figured out that these objects represent different stages in the lives of stars.



Three generations

The Lives of Stars

Stars do not last forever. Each star is born, goes through its life cycle, and eventually dies. (Of course, stars are not really alive. The words born, live, and die are just helpful comparisons.)

A Star is Born All stars begin their lives as parts of nebulas. A nebula is a large cloud of gas and dust spread out in an immense volume. A star, on the other hand, is made up of a large amount of gas in a relatively small volume.

In the densest part of a nebula, gravity pulls gas and dust together. A contracting cloud of gas and dust with enough mass to form a star is called a protostar. Proto means "earliest" in Greek, so a protostar is the earliest stage of a star's life.

A star is born when the contracting gas and dust from a nebula become so dense and hot that nuclear fusion starts. Recall that nuclear fusion is the process by which atoms combine to form heavier atoms. In the sun, for example, hydrogen atoms combine to form helium. During nuclear fusion, enormous amounts of energy are released. Nuclear fusion has not yet begun in a protostar.

Lifetimes of Stars How long a star lives depends on its mass. You might think that stars with more mass would last longer than stars with less mass. But the reverse is true. You can think of stars as being like cars. A small car has a small gas tank, but it also has a small engine that burns gas slowly. A large car has a larger gas tank, but it also has a larger engine that burns gas rapidly. So the small car can travel farther on a tank of gas than the larger car. Small-mass stars use up their fuel more slowly than large-mass stars, so they have much longer lives.

Generally, stars that have less mass than the sun use their fuel slowly, and can live for up to 200 billion years. Mediummass stars like the sun live for about 10 billion years. Astronomers think the sun is about 4.6 billion years old, so it is almost halfway through its lifetime.

Stars that have more mass than the sun have shorter lifetimes. A star that is 15 times as massive as the sun may live only about ten million years. That may seem like a long time, but it is only one tenth of one percent of the lifetime of the sun.

How long will a star that is the mass Checkpoint of the sun live?




FIGURE 12 **Young Stars**

New stars are forming in the nebula on top. The bottom photo shows a protostar in the Orion Nebula. Applying Concepts How do some of the gas and dust in a nebula become a protostar?






FIGURE 13 The Lives of Stars

A star's life history depends on its mass. A low-mass main-sequence star uses up its fuel slowly and eventually becomes a white dwarf. A high-mass star uses up its fuel quickly. After its supergiant stage, it will explode as a supernova, producing a neutron star or a black hole.

Interpreting Diagrams What type of star produces a planetary nebula?

Lab Skills Activity

Predicting

Find Algol, Sírius B, and Polaris in Figure 10, the H-R diagram. What type of star is each of these now? Predict what the next stage in each star's life will be.

Deaths of Stars


When a star begins to run out of fuel, its core shrinks and its outer portion expands. Depending on its mass, the star becomes either a red giant or a supergiant. All main-sequence stars eventually become red giants or supergiants. As shown in Figure 13, red giants and supergiants evolve in very different ways. After a star runs out of fuel, it becomes a white dwarf, a neutron star, or a black hole.

White Dwarfs Low-mass stars and medium-mass stars like the sun take billions of years to use up their nuclear fuel. As they start to run out of fuel, their outer layers expand, and they become red giants. Eventually, the outer parts grow larger still and drift out into space, forming a glowing cloud of gas called a planetary nebula. The blue-white core of the star that is left behind cools and becomes a white dwarf.

White dwarfs are only about the size of Earth, but they have about as much mass as the sun. Since a white dwarf has the same mass as the sun but only one millionth the volume, it is one million times as dense as the sun. A spoonful of material from a white dwarf has as much mass as a large truck. White dwarfs have no fuel, but they glow faintly from leftover energy. After billions of years, a white dwarf eventually stops glowing. Then it is called a black dwarf.



What is a white dwarf?



Supernovas The life cycle of a high-mass star is quite different from the life cycle of a low-mass or medium-mass star. High-mass stars quickly evolve into brilliant supergiants. When a supergiant runs out of fuel, it can explode suddenly. Within hours, the star blazes millions of times brighter. The explosion is called a **supernova**. After a supernova, some of the material from the star expands into space. This material may become part of a nebula. This nebula can then contract to form a new, partly recycled star. Astronomers think the sun began as a nebula that contained material from a supernova.

Neutron Stars After a supergiant explodes, some of the material from the star is left behind. This material may form a neutron star. **Neutron stars** are the remains of high-mass stars. They are even smaller and denser than white dwarfs. A neutron star may contain as much as three times the mass of the sun but be only about 25 kilometers in diameter, the size of a city.

In 1967, Jocelyn Bell, a British astronomy student, detected an object in space that appeared to give off regular pulses of radio waves. Some astronomers hypothesized that the pulses might be a signal from an extraterrestrial civilization. At first, astronomers even named the source LGM, for the "Little Green Men" in early science-fiction stories. Soon, however, astronomers concluded that the source of the radio waves was really a rapidly spinning neutron star. Spinning neutron stars are called **pulsars**, short for pulsating radio sources. Some pulsars spin hundreds of times per second!



Web Code: cfp-5043




FIGURE 14 Black Holes

The remains of the most massive stars collapse into black holes. This artist's impression shows a black hole pulling matter from a companion star. The material glows as it is pulled into the black hole. Applying Concepts If it is impossible to detect a black hole directly, how do astronomers find them?

Black Holes The most massive stars—those having more than 40 times the mass of the sun—may become black holes when they die. A black hole is an object with gravity so strong that nothing, not even light, can escape. After a very massive star dies in a supernova explosion, more than five times the mass of the sun may be left. The gravity of this mass is so strong that the gas is pulled inward, packing the gas into a smaller and smaller space. The gas becomes so densely packed that its intense gravity will not allow even light to escape. The remains of the star have become a black hole.

No light, radio waves, or any other form of radiation can ever get out of a black hole, so it is not possible to detect a black hole directly. But astronomers can detect black holes indirectly. For example, gas near a black hole is pulled so strongly that it revolves faster and faster around the black hole. Friction heats the gas up. Astronomers can detect X-rays coming from the hot gas and infer that a black hole is present. Similarly, if another star is near a black hole, astronomers can calculate the mass of the black hole from the effect of its gravity on the star. Scientists have detected dozens of star-size black holes with the Chandra X-ray Observatory. They have also detected huge black holes that are millions or billions of times the sun's mass.



What is a black hole?

Section / 3 Assessment

Target Reading Skill Sequencing Refer to your flowchart as you answer the questions.

Reviewing Key Concepts

- 1. a. Defining What is a nebula?
 - **b. Explaining** How does a star form from a nebula?
 - c. Comparing and Contrasting How is a protostar different from a star?
- 2. a. Identifying What factor determines how long a star lives?
 - b. Applying Concepts A star is twice as massive as the sun. Will its lifespan be longer, shorter, or the same as that of the sun?
- **3. a. Comparing and Contrasting** What is a white dwarf? How is it different from a neutron star?

- b. Relating Cause and Effect Why do some stars become white dwarfs and others become neutron stars or black holes?
- c. Predicting What will happen to the sun when it runs out of fuel? Explain.

Writing in Science

Descriptive Paragraph Write a description of one of the stages in the life of a star, such as a nebula, red giant, supernova, or white dwarf. Include information on how it formed and what will happen next in the star's evolution.

Star Systems and Galaxies

Reading Preview

Key Concepts

- What is a star system?
- What are the major types of galaxies?
- How do astronomers describe the scale of the universe?

Key Terms

- binary star
- eclipsing binary open cluster
- globular cluster galaxy
- spiral galaxy elliptical galaxy
- irregular galaxy quasar
- universe
 scientific notation

Target Reading Skill Building Vocabulary Carefully read the definition of each key

read the definition of each key term. Also read the neighboring sentences. Then write a definition of each key term in your own words.

Tab Discover Activity

Why Does the Milky Way Look Hazy?


- 1. Using a pencil, carefully poke at least 20 holes close together in a sheet of white paper.
- 2. Tape the paper to a chalkboard or dark-colored wall.
- 3. Go to the other side of the room and look at the paper. From the far side of the room, what do the dots look like? Can you see individual dots?

Think It Over

Making Models How is looking at the paper from the far side of the room like trying to see many very distant stars that are close together? How does your model compare to the photograph of the Milky Way below?

On a clear, dark night in the country, you can see a hazy band of light stretched across the sky. This band of stars is called the Milky Way. It looks as if the Milky Way is very far away. Actually, though, Earth is inside the Milky Way! The Milky Way looks milky or hazy from Earth because the stars are too close together for your eyes to see them individually. The dark blotches in the Milky Way are clouds of dust that block light from stars behind them.





Invisible Partners
If you saw someone dancing but
couldn't see a partner, you could
infer that the partner was there by
watching the dancer you could see.
Astronomers use a similar method

to detect faint stars in star systems.

Star Systems and Clusters


Our solar system has only one star, the sun. But this is not the most common situation for stars. Most stars are members of groups of two or more stars, called star systems. If you were on a planet in one of these star systems, at times you might see two or more suns in the sky! At other times, one or more of these suns would be below the horizon.

Multiple Star Systems Star systems that have two stars are called double stars or binary stars. (The prefix bi means "two.") Those with three stars are called triple stars. The nearby star Proxima Centauri may be part of a triple star system. The other two stars in the system, Alpha Centauri A and Alpha Centauri B, form a double star. Scientists are not sure whether Proxima Centauri is really part of the system or is just passing close to the other two stars temporarily.

Often one star in a binary star is much brighter and more massive than the other. Astronomers can sometimes detect a binary star even if only one of the stars can be seen from Earth. Astronomers can often tell that there is a dim star in a binary system by observing the effects of its gravity. As the dim companion star revolves around a bright star, the dim star's gravity causes the bright star to wobble back and forth. Imagine watching a pair of dancers who are twirling each other around. Even if one dancer were invisible, you could tell that the invisible dancer was there from watching the motion of the visible dancer.

Eclipsing Binaries A wobble is not the only clue that a star has a dim companion. A dim star in a binary star may pass in front of a brighter star and eclipse it. From Earth, the binary star would suddenly look much dimmer. A system in which one star periodically blocks the light from another is called an eclipsing binary. As Figure 16 shows, the star Algol is actually an eclipsing binary star system.

Eclipsing Binary Algol is an eclipsing binary star system consisting of a bright star and a dim companion. Each time the dimmer star passes in front of the brighter one, Algol appears less bright. Interpreting Diagrams When does Algol appear brighter?



Planets Around Other Stars In 1995, astronomers first discovered a planet revolving around another ordinary star. They used a method similar to the one used in studying binary stars. The astronomers observed that a star was moving slightly toward and away from us. They knew that the invisible object causing the movement didn't have enough mass to be a star. They inferred that it must be a planet.

Since then, astronomers have discovered more than 100 planets around other stars, and new ones are being discovered all of the time. Most of these new planets are very large, with at least half of the mass of Jupiter. A small planet would be hard to detect because it would have little gravitational effect on the star it orbited.

Could there be life on planets in other solar systems? Some scientists think it is possible. A few astronomers are using radio telescopes to search for signals that could not have come from natural sources. Such a signal might be evidence that an extraterrestrial civilization was sending out radio waves.

Star Clusters Many stars belong to larger groupings called star clusters. All of the stars in a particular cluster formed from the same nebula at about the same time and are about the same distance from Earth.

There are two major types of star clusters: open clusters and globular clusters. Open clusters have a loose, disorganized appearance and contain no more than a few thousand stars. They often contain many bright supergiants and much gas and dust. In contrast, globular clusters are large groupings of older stars. Globular clusters are round and densely packed with stars—some may contain more than a million stars.



What is a globular cluster?




FIGURE 17
Star Clusters
The stars in a globular cluster
(above) are all about the same age
and the same distance from Earth.
The Pleiades (left), also called the
Seven Sisters, is an open cluster.








FIGURE 18 Types of Galaxies There are three major types of galaxies: spiral, elliptical, and irregular.



For: Links on galaxies Visit: www.SciLinks.org Web Code: scn-0644

Galaxies

A galaxy is a huge group of single stars, star systems, star clusters, dust, and gas bound together by gravity. There are billions of galaxies in the universe. The largest galaxies have more than a trillion stars. Astronomers classify most galaxies into the following types: spiral, elliptical, and irregular. Figure 18 shows examples of these three.

Spiral Galaxies Some galaxies appear to have a bulge in the middle and arms that spiral outward, like pinwheels. Such galaxies are called spiral galaxies. The spiral arms contain many bright, young stars as well as gas and dust. Most new stars in spiral galaxies form in these spiral arms. Relatively few new stars are forming in the central bulge. Some spiral galaxies, called barred-spiral galaxies, have a huge bar-shaped region of stars and gas that passes through their center.


Elliptical Galaxies Not all galaxies have spiral arms. Elliptical galaxies look like round or flattened balls. These galaxies contain billions of stars but have little gas and dust between the stars. Because there is little gas or dust, stars are no longer forming. Most elliptical galaxies contain only old stars.

irregular Galaxies Some galaxies do not have regular shapes. These are known as irregular galaxies. Irregular galaxies are typically smaller than other types of galaxies. They generally have many bright, young stars and lots of gas and dust to form new stars.

Quasars In the 1960s, astronomers discovered objects that are very bright, but also very far away. Many of these objects are 10 billion light-years or more away, making them among the most distant objects in the universe. These distant, enormously bright objects looked almost like stars. Since quasi means "something like" in Latin, these objects were given the name quasi-stellar objects, or quasars.

What could be so bright at such a great distance from Earth? Astronomers have concluded that quasars are active young galaxies with giant black holes at their centers. Each of these black holes has a mass a billion times or more as great as that of the sun. As enormous amounts of gas revolve around the black hole, the gas heats up and shines brightly.








FIGURE 19
Structure of the Milky Way
From the side, the Milky Way
appears to be a narrow disk with a
bulge in the middle. The galaxy's
spiral structure is visible only from
above or below the galaxy.
Interpreting Diagrams Where in
the galaxy is the sun located?

The Wilky Way

Our solar system is located in a spiral galaxy called the Milky Way. As Figure 19 shows, the shape of the Milky Way varies depending on your vantage point. From the side, the Milky Way would look like a narrow disk with a large bulge in the middle. But from the top or bottom, the Milky Way would have a spiral, pinwheel shape. You can't see the spiral shape of the Milky Way from Earth because our solar system is inside the galaxy in one of the spiral arms.

The Milky Way is usually thought of as a standard spiral galaxy. However, recent evidence suggests that the Milky Way is a barred-spiral galaxy instead.

When you see the Milky Way at night during the summer, you are looking toward the center of our galaxy. The center of the galaxy is about 25,000 light-years away, but it is hidden from view by large clouds of dust and gas. However, astronomers can study the center using X-rays, infrared radiation, and radio waves.

Reading Checkpoint

How far away is the center of the galaxy?


Lab Try This Activity

A Spiral Galaxy

You can make a model of our galaxy.

- Using pipe cleaners, make a pinwheel with two spirals.
- 2. View the spirals along the surface of the table.
 Sketch what you see.
- 3. Next, view the spirals from above the table and sketch them.

Observing The sun is inside a flat spiral galaxy. From Earth's position on the flat surface, is it possible to get a good view of stars in the spiral arms? Why or why not?



Math Skills

Scientific Notation

The bright star Deneb is about 3,230 light-years from Earth. To express this number in scientific notation, first insert a decimal point in the original number so that you have a number between one and ten. In this case, the number is 3.23.

To determine the power of 10, count the number of places that the decimal point moved. Here the decimal point moved three places.

3,230 light-years = 3.23×10^3 light-years


Practice Problem The sun takes about 220,000,000 years to revolve once around the center of the galaxy. Express this length of time in scientific notation.

The Scale of the Universe

Astronomers define the universe as all of space and everything in it. The universe is enormous, almost beyond imagination. Astronomers study objects as close as the moon and as far away as quasars. They study incredibly large objects, such as galaxies that are millions of light-years across. They also study the behavior of tiny particles, such as the atoms within stars. Since the numbers astronomers use are often very large or very small, they frequently use scientific notation to describe sizes and distances in the universe.

Scientific Notation Scientific notation uses powers of ten to write very large or very small numbers in shorter form. Each number is written as the product of a number between 1 and 10 and a power of 10. For example: 1,200 is written as 1.2×10^3 . One light-year is about 9,500,000,000,000 meters. Since there are 15 digits after the first digit, in scientific notation this number is written as 9.5×10^{15} meters.

The Immensity of Space The structures in the universe vary greatly in scale. To understand the scale of these structures, imagine that you are going on a journey through the universe. Refer to Figure 20 as you take your imaginary trip. Start at the left with something familiar—a girl looking through binoculars. She is about 1.5 meters tall. Now shift to the right and change the scale by 10,000,000 or 10^7 . You're now close to the diameter of Earth, 1.28×10^7 meters. As you move from left to right across Figure 20, the scale increases. The diameter of the sun is about 100 times that of Earth.



Beyond the solar system, the sizes of observable objects become much larger. For example, within our galaxy, the beautiful Cat's Eye Nebula is about 3×10^{16} meters across.

Beyond our galaxy are billions of other galaxies, many of which contain billions of stars. For example, the nearby spiral galaxy Andromeda is about 2×10^{21} meters across. The Milky Way is part of a cluster of 50 or so galaxies called the Local Group. The Local Group is part of the Virgo Supercluster, which contains hundreds of galaxies. The size of the observable universe is about 10^{10} light years, or 10^{26} meters.

FIGURE 20
Scientific Notation
Scientists often use scientific notation to help describe the vast sizes and distances in space.
Calculating About how many times larger is the Cat's Eye Nebula than Earth?

Section 4 Assessment

Target Reading Skill Building Vocabulary Use your definitions to help answer the questions.

Reviewing Key Concepts

- 1. a. Defining What is a binary star?
 - **b. Classifying** Are all binary stars part of star systems? Explain.
 - c. Applying Concepts Some binary stars are called eclipsing binaries. Explain why this term is appropriate. (*Hint:* Think about Algol as you write your answer.)
- 2. a. Listing Name the main types of galaxies.
 - **b.** Classifying What type of galaxy is the Milky Way?
 - c. Classifying Suppose astronomers discover a galaxy that contains only old stars. What type of galaxy is it likely to be?

- **3. a.** Reviewing What is scientific notation?
 - b. Explaining How is scientific notation useful to astronomers?
 - c. Calculating How large is the Cat's Eye Nebula in light-years? (*Hint:* Refer to Figure 20.)

Math Practice

4. Scientific Notation The star Betelgeuse has a diameter of 940,000,000 km. Betelgeuse is 427 light-years from Earth. Write each of these figures in scientific notation.

The Expanding Universe

Reading Preview

Key Concepts

- What is the big bang theory?
- How did the solar system form?
- What do astronomers predict about the future of the universe?

Key Terms

- big bang Hubble's law
- cosmic background radiation.
- solar nebula planetesimal
- dark matter dark energy

Zane Discover Activity

How Does the Universe Expand?


- 1. Use a marker to put 10 dots on an empty balloon. The dots represent galaxies.
- 2. Blow up the balloon. What happens to the distances between galaxies that are close together? Galaxies that are far apart?

Think It Over

Inferring If the universe is expanding, do galaxies that are close together move apart faster or slower than galaxies that are far apart? Explain.

Target Reading Skill Identifying Supporting

Evidence As you read, identify the evidence that supports the big bang theory. Write the evidence in a graphic organizer like the one below.



The Andromeda Galaxy is the most distant object that the human eye can see. Light from this galaxy has traveled for about 3 million years before reaching Earth. When that light finally reaches your eye, you are seeing how the galaxy looked 3 million years ago. It is as though you are looking back in time.

Astronomers have photographed galaxies that are billions of light-years away. Light from these galaxies traveled for billions of years before it reached Earth. From these observations, astronomers are able to infer the age of the universe.

How the Universe Formed

Astronomers theorize that the universe began billions of years ago. At that time, the part of the universe we can now see was no larger than the period at the end of this sentence. This tiny universe was incredibly hot and dense. The uni-

verse then exploded in what astronomers call the big bang.

Nearly every visible object in this image is a distant galaxy.



According to the big bang theory, the universe formed in an instant, billions of years ago, in an enormous explosion. Since the big bang, the size of the universe has been increasing rapidly. The universe is billions of times larger now than it was early in its history.

As the universe expanded, it gradually cooled. After a few hundred thousand years, atoms formed. About 200 million years after the big bang, the first stars and galaxies formed.

If the big bang theory is accurate, what evidence might you expect to find in today's universe? You might expect that the matter that had been hurled apart by the big bang would still be moving apart. You might also expect to find evidence of energy left over from the explosion.

Moving Galaxies An American astronomer, Edwin Hubble, discovered important evidence that later helped astronomers to develop the big bang theory. In the 1920s, Hubble studied the spectrums of many galaxies at various distances from Earth. By examining a galaxy's spectrum, Hubble could tell how fast the galaxy is moving and whether it is moving toward our galaxy or away from it.

Hubble discovered that, with the exception of a few nearby galaxies, all galaxies are moving away from us and from each other. Hubble found that there is a relationship between the distance to a galaxy and its speed. **Hubble's law** states that the farther away a galaxy is, the faster it is moving away from us. Hubble's law strongly supports the big bang theory.

FIGURE 21
Retreating Galaxies
All of the distant galaxies
astronomers have observed are
moving rapidly away from our
galaxy and from each other.

Math

Analyzing Data

Speeding Galaxies

Use the graph to answer the questions below about moving clusters of galaxies.

- 1. Reading Graphs How far away is the Bootes cluster? How fast is it moving?
- 2. Reading Graphs Which galaxy is moving away the fastest? Which galaxy is closest to Earth?
- 3. Drawing Conclusions How are the distance and speed of a galaxy related?
- 4. Predicting Predict the speed of a galaxy that is 5 billion light-years from Earth.





FIGURE 22 Rising Dough

The galaxies in the universe are like the raisins in rising bread dough. Making Models How does rising raisin bread dough resemble the expanding universe?



To understand how the galaxies are moving, think of raisin bread dough that is rising. If you could shrink yourself to sit on a raisin, you would see all the other raisins moving away from you. The farther a raisin was from you, the faster it would move away, because there would be more bread dough to expand between you and the raisin. No matter which raisin you sat on, all the other raisins would seem to be moving away from you. You could tell that the bread dough was expanding by watching the other raisins.

The universe is like the bread dough. Like the raisins in the dough, the galaxies in the universe are moving away from each other. In the universe, it is space that is expanding, like the dough between the raisins.

Cosmic Background Radiation In 1965, two American physicists, Arno Penzias and Robert Wilson, accidentally detected faint radiation on their radio telescope. This mysterious glow was coming from all directions in space. Scientists later concluded that this glow, now called cosmic background radiation, is the leftover thermal energy from the big bang. This energy was distributed in every direction as the universe expanded.

Age of the Universe Since astronomers can measure approximately how fast the universe is expanding now, they can infer how long it has been expanding. Based on careful measurements of how fast distant galaxies are moving away from us and the cosmic background radiation, astronomers estimate that the universe is about 13.7 billion years old.

Formation of the Solar System

After the big bang, matter in the universe separated into galaxies. Gas and dust spread throughout space. Where the solar system is now, there was only cold, dark gas and dust. How did the solar system form? The leading hypothesis is explained below.


The Solar Nebula About five billion years ago, a giant cloud of gas and dust collapsed to form our solar system. A large cloud of gas and dust such as the one that formed our solar system is called a solar nebula. Slowly, gravity began to pull the solar nebula together. As the solar nebula shrank, it spun faster and faster. The solar nebula flattened, forming a rotating disk. Gravity pulled most of the gas into the center of the disk, where the gas eventually became hot and dense enough for nuclear fusion to begin. The sun was born.


Planetesimals In the outer parts of the disk, gas and dust formed small asteroid-like and comet-like bodies called **planetesimals**. These formed the building blocks of the planets. Planetesimals collided and grew larger by sticking together, eventually combining to form the planets.


The Inner Planets When the solar system formed, temperatures were very high. It was so hot close to the sun that most water and other ice-forming materials simply vaporized. Most gases escaped the gravity of the planets that were forming in this region. As a result, the inner planets, Mercury, Venus, Earth, and Mars, are relatively small and rocky.

The Outer Planets In contrast, farther from the sun it was much cooler. As the planets in this region grew, their gravity increased and they were able to capture much of the hydrogen and helium gas in the surrounding space. As a result, the planets Jupiter, Saturn, Uranus, and Neptune became very large. Most comets formed near Jupiter and Saturn. They were later flung out to the outer solar system. Beyond the gas giants, a huge disk of ice and other substances formed. Pluto also formed in this region.













FIGURE 23
How the Solar System Formed
The solar system formed from a
collapsing cloud of gas and dust.



For: Links on the expanding universe Visit: www.SciLinks.org Web Code: scn-0645

FIGURE 24 Vera Rubin Astronomer Vera Rubin's observations proved the existence of dark matter.



The Future of the Universe

What will happen to the universe in the future? One possibility is that the universe will continue to expand, as it is doing now. All of the stars will eventually run out of fuel and burn out, and the universe will be cold and dark. Another possibility is that the force of gravity will begin to pull the galaxies back together. The result would be a reverse big bang, or "big crunch." All of the matter in the universe would be crushed into an enormous black hole.

Which of these possibilities is more likely? Recent discoveries have produced a surprising new view of the universe that is still not well understood. New observations lead many astronomers to conclude that the universe will likely expand forever.


Dark Matter Until fairly recently, astronomers assumed that the universe consisted solely of the matter they could observe directly. But this idea was disproved by the American astronomer Vera Rubin. Rubin made detailed observations of the rotation of spiral galaxies. She discovered that the matter that astronomers can see, such as stars and nebulas, makes up as little as ten percent of the mass in galaxies. The remaining mass exists in the form of dark matter.

Dark matter is matter that does not give off electromagnetic radiation. Dark matter cannot be seen directly. However, its presence can be inferred by observing the effect of its gravity on visible objects, such as stars, or on light.

Astronomers still don't know much about dark matter—what it is made of or all of the places where it is found. But astronomers estimate that about 23 percent of the universe's mass is made of dark matter.

An Accelerating Expansion In the late 1990s, astronomers observed that the expansion of the universe appears to be accelerating. That is, galaxies seem to be moving apart at a faster rate now than in the past. This observation was puzzling, as no known force could account for it. Astronomers infer that a mysterious new force, which they call dark energy, is causing the expansion of the universe to accelerate. Current estimates indicate that most of the universe is made of dark energy and dark matter.

Astronomy is one of the oldest sciences, but there are still many discoveries to be made and puzzles to be solved about this universe of ours!



What is the effect of dark energy?




FIGURE 25 Dark Matter

Astronomers measured the effect of gravity on light to produce this computer image of how dark matter (in blue) is distributed across a cluster of galaxies.

Section 5 Assessment

Target Reading Skill Identifying Supporting Evidence Refer to your graphic organizer about the big bang theory as you answer Question 1 below.

Reviewing Key Concepts

- 1. a. Defining What was the big bang?
 - **b.** Summarizing When did the big bang occur?
 - c. Describing Describe two pieces of evidence that support the big bang theory.
- 2. a. Summarizing How old is the solar system?
 - **b.** Relating Cause and Effect What force caused the solar system to form?
 - c. Sequencing Place the following events in the proper order: planets form; planetesimals form; solar nebula shrinks; nuclear fusion begins in the sun.

- 3. a. Defining What is dark matter?
 - **b. Explaining** How do scientists know that dark matter exists?
 - c. Predicting What evidence has led scientists to predict that the universe will continue to expand forever?

Lab At-Home Activity

Stargazing Plan an evening of stargazing with adult family members. Choose a dark, clear night. Use binoculars if available and the star charts in the appendix to locate the Milky Way and some interesting stars that you have learned about. Explain to your family what you know about the Milky Way and each constellation that you observe.

Chapter /A

Study Guide

The BIO Islea

Structure of the Universe Astronomers learn about the structure of the universe and how it has changed over time by studying stars, galaxies, and other objects in space

1 Telescopes

Key Concepts

- The electromagnetic spectrum includes radio waves, infrared radiation, visible light, ultraviolet radiation, X-rays, and gamma rays.
- Telescopes are instruments that collect and focus light and other forms of electromagnetic radiation.
- Many large observatories are located on mountaintops or in space.

Key Terms

- telescope
- visible light
- wavelength
- spectrum
- optical telescope
- electromagnetic radiation
- refracting telescope
- convex lens
- reflecting telescope
- radio telescope
- observatory

2 Characteristics of Stars

Key Concepts

- Characteristics used to classify stars include color, temperature, size, composition, and brightness.
- The brightness of a star depends upon both its size and temperature.
- Astronomers use a unit called the light-year to measure distances between the stars.
- Astronomers often use parallax to measure distances to nearby stars.
- Astronomers use H-R diagrams to classify stars and to understand how stars change over time.

Key Terms

- constellation spectrograph
- apparent brightness
 absolute brightness
- light-year parallax
- Hertzsprung-Russell diagram
- main sequence

3 Lives of Stars

Key Concepts

- A star is born when nuclear fusion starts.
- How long a star lives depends on its mass.
- After a star runs out of fuel, it becomes a white dwarf, a neutron star, or a black hole.

Key Terms

- nebula protostar white dwarf
- supernova neutron star pulsar
- black hole

4 Star Systems and Galaxies

Key Concepts

- Most stars are members of groups of two or more stars called star systems.
- Astronomers classify most galaxies into the following types: spiral, elliptical, and irregular.
- Our solar system is located in a spiral galaxy called the Milky Way.
- Astronomers often use scientific notation to describe sizes and distances in the universe.

Key Terms

- binary star eclipsing binary open cluster
- globular cluster galaxy spiral galaxy
- elliptical galaxy irregular galaxy
- quasar universe scientific notation

5 The Expanding Universe

Key Concepts

- According to the big bang theory, the universe formed in an instant, billions of years ago, in an enormous explosion.
- About five billion years ago, a giant cloud of gas and dust collapsed to form our solar system.
- New observations lead astronomers to conclude that the universe will likely expand forever.

Key Terms

- big bang Hubble's law
- cosmic background radiation solar nebula
- planetesimal
 dark matter
 dark energy

Review and Assessment



For: Self-Assessment Visit: PHSchool.com Web Code: cfa-5040

Organizing Information

Concept Mapping Copy the concept map about telescopes onto a separate sheet of paper. Then complete it and add a title. (For more on Concept Mapping, see the Skills Handbook.)

Telescopes

nto

can be

r
the

On Earth
the

collect types of radiation

b. ?

Visible light

c. ?

Gamma rays

Reviewing Key Terms

Choose the letter of the best answer.

- 1. Visible light is a form of
 - a. spectrum.
 - b. electromagnetic radiation.
 - c. wavelength.
 - d. cosmic background radiation.
- 2. An H-R diagram is a graph of stars' temperature and
 - a. apparent brightness.
 - b. main sequence.
 - c. absolute brightness.
 - d. parallax.
- 3. A low-mass main sequence star will eventually evolve into a
 - a. white dwarf.
- **b.** protostar.
- c. black hole.
- d. nebula.
- 4. A star system in which one star blocks the light from another is called a(n)
 - a. open cluster.
 - b. quasar.
 - c. binary star.
 - d. eclipsing binary.
- 5. Astronomers theorize that the universe began in an enormous explosion called the
 - a. solar nebula.
 - b. supernova.
 - c. big bang.
 - d. big crunch.

If the statement is true, write *true*. If it is false, change the underlined word or words to make the statement true.

- **6.** A <u>reflecting telescope</u> uses convex lenses to gather and focus light.
- 7. Astronomers use <u>spectrographs</u> to determine the chemical composition of stars.
- 8. Pulsars are a kind of neutron star.
- A galaxy shaped like a ball and containing only older stars is most likely a <u>spiral galaxy</u>.
- **10.** Globular clusters are small asteroid-like bodies that formed the building blocks of the planets.

Writing in Science

News Article Imagine that you are a journalist covering current research in astronomy, including stars and black holes. Write an article explaining what black holes are, how they form, and how they can be detected.



Stars, Galaxies, and the Universe
Video Preview
Video Field Trip
Video Assessment

Review and Assessment

Checking Concepts


- 11. Is a light-year a unit of distance or a unit of time? Explain.
- 12. Why can't astronomers measure the parallax of a star that is a million light-years away?
- 13. At what point in the evolution of a star is the star actually born?
- **14.** Where in our galaxy does most star formation take place?
- 15. What is Hubble's law?
- **16.** How can astronomers detect dark matter if they cannot observe it directly?

Math Practice

- 17. Calculating The bright star Spica is 262 lightyears from our solar system. How many kilometers is this?
- 18. Scientific Notation The star Antares is approximately 604 light-years from Earth. Write this distance in scientific notation.

Thinking Critically

19. Inferring What advantage might there be to locating a telescope, such as the one shown below, on the moon?




20. Applying Concepts Describe a real-world situation involving absolute and apparent brightness. (*Hint:* Think about riding in a car at night.)

- 21. Relating Cause and Effect How does a star's mass affect its lifetime?
- 22. Comparing and Contrasting Compare the conditions that led to the formation of the terrestrial planets with those that led to the formation of the gas giants.

Applying Skills

Use the data in the H-R diagram below to answer Questions 23–26.

Hertzsprung-Russell Diagram



- 23. Interpreting Diagrams Which star has a greater absolute brightness, Aldebaran or Sirius B?
- 24. Interpreting Diagrams Which stars have higher surface temperatures than Sirius A?
- 25. Applying Concepts Which star is most likely to be red: Rigel, Sirius B, or Betelgeuse?
- **26.** Comparing and Contrasting Compare Aldebaran and the sun in terms of size, temperature, and absolute brightness.

Zone Chapter Project

Performance Assessment Check the final draft of your constellation story for correct spelling, grammar, punctuation, and usage. Then decide how you will present your story. For example, you could make a poster, read your story aloud, or perform it as a skit or a play.

Standardized Test Prep

Test-Taking Tip

Sequencing Events

Some questions ask you to arrange a series of events in order. For example, you might be asked which event comes first or last, or which event comes before another event. Before looking at the answer choices, first try to recall the sequence of events in the entire process. If you have an idea of the sequence beforehand, you should find it easier to identify the correct answer.

Sample Question

Which of the following correctly describes the evolution of a sun-like star from young to old?

- A white dwarf, red giant, main-sequence star, protostar
- **B** red giant, main-sequence star, white dwarf, protostar
- C protostar, main-sequence star, white dwarf, red giant
- **D** protostar, main-sequence star, red giant, white dwarf

Answer

The correct answer is **D**. Choice **A** gives the correct order from old to young, rather than young to old. Choices **B** and **C** do not correctly sequence the life cycle of any star.

Choose the letter of the best answer.

- 1. The most common chemical element in most stars is
 - A oxygen.
 - **B** hydrogen.
 - C helium.
 - D nitrogen.
- 2. The main factor that affects the evolution of a star is its
 - F color.
 - G apparent brightness.
 - H mass.
 - J parallax.

- 3. The color of a star is related to its temperature. Which of the following color sequences correctly identifies the temperatures of stars in order from hottest to coldest?
 - A red, red-orange, yellow, white, blue
 - B yellow, white, blue, red, red-orange
 - C blue, yellow, red-orange, red, white
 - D blue, white, yellow, red-orange, red

The table below gives an estimate of the distribution of stars in the Milky Way galaxy. Use the table and your knowledge of science to answer Questions 4 and 5.

Type of Star	Percentage of Total		
Main sequence	90.75%		
Red Giant	0.50%		
Supergiant	< 0.0001%		
White Dwarf	8.75%		

- 4. According to the table, the most common type of stars in the Milky Way is
 - F main-sequence stars.
 - G red giants.
 - H supergiants.
 - J white dwarfs.
- 5. If there are a total of 400 billion stars in the Milky Way, about how many white dwarfs are there in the galaxy?
 - A 8.75 billion
 - B 35 billion
 - C 87.5 billion
 - D 3,500 billion

Constructed Response

6. Describe the appearance of the Milky Way as you would see it both from Earth and from a point directly above or below the galaxy. Why does the galaxy look different from different vantage points?

Interdisciplinary **Exploration**

Journey to Mars

The six-wheeled rover inched onto the surface of Mars.

Scientists on Earth held their breaths.

Then, Spirit hummed into action.

Spirit was the first star of the 2004 Mars mission. Engineers at the Jet Propulsion Laboratory in Pasadena, California, guided the rover from Earth by remote control. Spirit carried a high-tech microscope, cameras, and geologic instruments. Within hours of landing in Gusev Crater, Spirit was beaming images of the rocks and red soil back to Earth. Engineers on Earth "drove" Spirit to its first target, a large rock that they named "Adirondack."

A major goal of the Mars Exploration Mission was to look for evidence of past liquid water on Mars. Earlier photos of the red planet lead scientists to believe that Gusev Crater was once a dried-up lake bed. The presence of water increases the likelihood that life may have once existed on Mars.

Just three weeks later, another rover called *Opportunity* landed on the opposite side of Mars in a strange, flat landscape. It sent back images of a shallow red crater with bedrock in the distance.

